
Optimal Bt Toxins and Gene Silencing RNAs for Management of ACP to Mitigate the Impact of HLB

Researchers: Bryony Bonning, Nabil Killiny, Chooa El-Mohtar, Manjul Dutt, Lukasz Stelinski, Monique Rivera

Contact: Bryony Bonning

bbonning@ufl.edu,

UF/IFAS Gainesville or

Nabil Killiny

nabilkilliny@ufl.edu

UF/IFAS CREC



Currently, HLB is managed primarily by using pesticides to control the vector of HLB, the Asian citrus psyllid (ACP). Traditional pesticides are expensive for growers, harmful to beneficial insects, and may contaminate the environment. In addition, ACP is becoming resistant to many of the most widely used pesticides, rendering them ineffective. Therefore, new approaches to managing ACP are sought. In this project, we aim to combine RNA interference technology (RNAi) with *Bacillus thuringiensis* (Bt) bioinsecticidal proteins to gain

a new and efficient method of control for ACP. The pesticidal proteins produced by Bt have been widely used for crop protection in agricultural commodities such as corn and cotton for years without harm to humans, domestic animals, or beneficial insects. However, its use in ACP control has been challenging. RNAi is the application of exogenous double-stranded RNAs (dsRNA), which will block targeted genes such as detoxification enzymes or genes controlling immune response, to reduce their activity in the insect. Genes targeting sucrose hydrolase

and aquaporin for disrupting the sugar and water metabolism of ACP are currently being tested. The use of both RNAi and Bt pesticidal proteins in combination may be more effective in reducing ACP survival, than use of either technology alone. The study of different combinations of RNAi gene targets and Bt proteins will reveal the best combinations for ACP control, and provide growers with an eco-friendly alternative to pesticide use for psyllid management.

Funding

 National Institute of Food and Agriculture
U.S. DEPARTMENT OF AGRICULTURE