

Table 1. Germplasm in Aurantiodeae subfamily^a tested for resistance to *Diaprepes abbreviatus*.

Tribe	Subtribe	Group	Species	Ref. ^b
Clauseneae	Micromelinae		None tested	---
Clauseneae	Clauseninae		<i>Murraya paniculata</i> (L.) Jack	4
Clauseneae	Clauseninae		<i>Glycosmis pentaphylla</i> (Retz.) Correa	3, 4
Clauseneae	Merrilliinae		None tested	---
Citreae	Triphasiinae		<i>Triphasia trifolia</i> (Burm. f.) P. Wils.	4
Citreae	Citrinae	A	<i>Severinia buxifolia</i> (Poir.) Tenore	1, 4
Citreae	Citrinae	B	<i>Citropsis gilleuana</i> Swing. & M. Kell.	4
Citreae	Citrinae	C	<i>Fortunella hindsii</i> (Champ.) Swing.	4
Citreae	Citrinae	C	<i>Eremocitrus glauca</i> (Lindl.) Swing.	4
Citreae	Citrinae	C	<i>Poncirus trifoliata</i> (L.) Raf.	1
Citreae	Citrinae	C	<i>Clymenia polyandra</i> (Tan.) Swing.	4
Citreae	Citrinae	C	<i>Microcitrus australasica</i> (F. Muell.) Swing.	1
Citreae	Citrinae	C	<i>Microcitrus australis</i> (Planch.) Swing.	4
Citreae	Citrinae	C	<i>Citrus aurantium</i> L.	1, 2, 4
Citreae	Citrinae	C	<i>Citrus hystrix</i> DC.	4
Citreae	Citrinae	C	<i>Citrus limon</i> (L.) Burm. f.	1
Citreae	Citrinae	C	<i>Citrus macrophylla</i> Webster	1, 2
Citreae	Citrinae	C	<i>Citrus reticulata</i> Blanco	1, 2, 4
Citreae	Citrinae	C	<i>Citrus sinensis</i> (L.) Osbeck	1
Citreae	Basamocitrinae	A	None tested	---
Citreae	Basamocitrinae	B	<i>Afraegle paniculata</i> (Schum.) Engl.	4
Citreae	Basamocitrinae	B	<i>Balsamocitrus dawei</i> Stapf.	4
Citreae	Basamocitrinae	C	None tested	---

^aSwingle (1967).

^b1 = Beavers and Hutchison (1985); 2 = Shapiro and Gottwald (1995); 3 = Shapiro et al. (1997); 4 = This manuscript.

Table 2. Common and new hybrid rootstocks tested for resistance to *Diaprepes abbreviatus*.

Name/Code	Parentage	Commercial utilization ^a
Carrizo citrange	<i>Citrus sinensis</i> x <i>P. trifoliata</i>	24
Swingle citrumelo	<i>C. paradisi</i> Macf. x <i>P. trifoliata</i>	53
Smooth Flat Seville	<i>Citrus</i> hybrid, parentage unknown	4
Sun Chu Sha	<i>C. reticulata</i>	2
Sour orange	<i>C. aurantium</i>	2
Yuma	probably <i>C. paradisi</i> x <i>P. trifoliata</i>	0
HRS-801	<i>C. reticulata</i> x <i>P. trifoliata</i>	0
HRS-802	<i>C. grandis</i> (L.) Osbeck x <i>P. trifoliata</i>	0
HRS-812	<i>C. reticulata</i> x <i>P. trifoliata</i>	0
HRS-896	<i>C. reticulata</i> x <i>P. trifoliata</i>	0
HRS-941	<i>C. reticulata</i> x <i>P. trifoliata</i>	0
HRS-1101	<i>C. reticulata</i> x <i>C. macrophylla</i>	0

^aPercentage of Florida citrus nursery trees propagated on this rootstock over the last 7 years, as indicated by Florida Dept. of Agriculture (1998).




Fig. 1. Relationship of noninfested root weight to infested root consumed.




Fig. 2. Relationship of noninfested root weight to larval weight gain.

to allow larvae to feed on the roots. During this initial 6 week period, *D. abbreviatus* larvae feed and grow most rapidly. After this feeding period, roots were washed free of soil mix and recovered larvae were counted and weighed (Shapiro et al., 1997). Root masses of both infested and control plants were severed at the soil line, air-dried, and weighed. The average root weight of seven noninfested plants of each genotype was calculated at the end of the experiment. The amount of root loss due to feeding of *D. abbreviatus* was estimated for each replicate (pot) as the difference between this mean value and the root weight for each infested plant. A comparison of infested to noninfested plants of each cultivar independently by *t* test was used to determine resistance based on root damage because of uncontrolled factors that varied between and within experiments. Shapiro and Gottwald (1995) expressed root loss as a ratio of infested root weight : uninfested root weight. We did not use this conversion because our initial root weights varied considerably and a ratio comparison would bias resistance estimates to favor selections with larger initial root weights. Instead, resistance to root damage by larvae was assessed by a *t* test comparison of root weight for infested and noninfested plants of each selection. Resistant types were those with no significant difference ($P \leq 0.05$) in root dry weight between treated plants and the noninfested controls.

For comparison of means for larval growth and survival among rootstocks and species within experiments, a single factor analysis of variance (ANOVA) was used, followed by Tukey's honestly significant difference (HSD) test (SAS version 6.08, SAS Institute; and Statistica version 5.0, Statsoft, Tulsa, Okla.). All statistical comparisons used 'Carrizo' citrange as the standard because it is a commonly planted rootstock and is widely accepted as susceptible to *D. abbreviatus*.

For easy comparisons of cultivars/selections included in different experiments, root

BREEDING, CULTIVARS, ROOTSTOCKS, & GERMPLASM RESOURCES

loss and larval growth were converted to a numerical value that ranged from 1 to 4, where 1 = highly susceptible, 2 = susceptible (by definition = 'Carrizo'), 3 = resistant, and 4 = highly resistant. For the nine selections tested in more than one experiment, the summary rating of each of the resistance measures was expressed as an average of the values from each experiment. An overall resistance rating (average of the measures of root loss and larval growth) was also calcu-

lated. This rating is proposed as an aid for quantifying levels of resistance in a preliminary study until more detailed information can be collected on resistance of individual selections.

Results

Each experiment identified one or more citrus rootstocks or citrus relatives that differed significantly from 'Carrizo' by at least

one of the measures of resistance studied (Tables 3 and 4). Both highly susceptible and resistant types were identified within some individual tests, and results for genotypes included in more than one test were consistent from one test to another for most cultivars.

Shapiro and Gottwald (1995) suggested that the rootstock 'Swingle' citrumelo was somewhat more resistant to the weevil than 'Carrizo'. However, during repeated testing we found no consistent difference between the two cultivars. In particular, root loss, larval survival, and larval growth were statistically indistinguishable for the two cultivars in five of the six experiments (Tables 3 and 4). In the remaining experiment (Expt. 1), less larval growth occurred on 'Swingle' than on 'Carrizo', while root loss and larval survival were indistinguishable.

Because 'Carrizo' and 'Swingle' appeared similar in weevil resistance, data collected for the two cultivars were used to examine the relationship between root loss from infestation and noninfested root weight (Fig. 1). There was a correlation between these two parameters for both 'Carrizo' ($r^2 = 0.726, P < 0.031$) and 'Swingle' ($r^2 = 0.674, P < 0.045$).

In three of the six experiments, no significant differences in percentage root loss were observed between the cultivars/species tested. Plants of *B. dawei*, *M. paniculata*, *C. gillettiana*, 'Smooth Flat Seville', 'Sun Chu Sha', *G. pentaphylla*, and the hybrid HRS-802 did not suffer significant root loss from feeding of *D. abbreviatus* in at least one experiment.

The relationship between noninfested root weight and larval weight gain was nonsignificant for either 'Carrizo' ($r^2 = 0.056, P < 0.651$) or 'Swingle' ($r^2 = 0.319, P < 0.243$) (Fig. 2). We conclude that larval growth and survival were not limited by availability of food, demonstrating that our level of infestations and test duration were appropriate to minimize competition between larvae for nutritional resources. To evaluate resistance by larval growth, the larval weight gained on each species was compared with 'Carrizo' for each experiment. Several of the species supported significantly less larval growth than 'Carrizo', including *B. dawei*, *C. polyandra*, *S. buxifolia*, *M. australis*, *G. pentaphylla*, *E. glauca*, *T. trifolia*, *C. hystris*, and hybrids HRS-801, HRS-896, and HRS-941 (Table 4).

Some cultivars/species differed significantly from 'Carrizo' in larval survival in some experiments. However, larval survival appeared to follow the same patterns as larval growth, but was somewhat less sensitive than the latter measure of plant resistance. All significant differences that were observed in larval survival were mirrored by significant differences in larval growth. Therefore, larval survival was not included in the overall summary and resistance rating described below. Many of the 23 other genotypes tested during the course of the six experiments differed little from the standard, 'Carrizo', in apparent susceptibility to *D. abbreviatus* larvae (Tables 3 and 4).

Table 3. Mean root damage to rootstocks in six experiments, means \pm SE.¹

Species/cultivar	Root wt (g)		P ²	Rating ³
	Noninfested	Infested		
<i>Expt. 1. Jan. 1996</i>				
Carrizo	34.9 \pm 3.0	14.1 \pm 1.2	0.000	S
Swingle	18.3 \pm 1.8	7.6 \pm 1.2	0.000	S
Sun Chu Sha	47.2 \pm 5.0	21.8 \pm 3.6	0.001	S
HRS-801	16.0 \pm 0.9	8.0 \pm 0.7	0.000	S
HRS-802	75.2 \pm 4.6	39.8 \pm 3.3	0.000	S
HRS-812	33.5 \pm 4.0	12.4 \pm 1.5	0.000	S
HRS-896	30.7 \pm 3.8	13.1 \pm 1.1	0.001	S
HRS-941	29.3 \pm 2.6	10.7 \pm 1.2	0.000	S
<i>Afraegle paniculata</i> ⁴	57.7 \pm 2.9	30.6 \pm 4.8	0.000	S
<i>Expt. 2. June 1996</i>				
Carrizo	44.6 \pm 8.3	21.3 \pm 1.6	0.018	S
Swingle	37.1 \pm 1.7	13.1 \pm 1.4	0.000	S
HRS-801	23.1 \pm 3.8	7.1 \pm 1.5	0.002	S
HRS-802	39.5 \pm 6.3	35.9 \pm 5.0	0.655	R
HRS-896	24.4 \pm 2.1	7.9 \pm 1.4	0.000	S
Smooth Flat Seville	64.5 \pm 20.6	52.2 \pm 6.0	0.576	R
<i>Balsamocitrus dawei</i>	42.0 \pm 8.6	46.0 \pm 12.9	0.802	R
Yuma	55.4 \pm 4.9	34.8 \pm 3.3	0.005	S
<i>Citropsis gillettiana</i>	72.9 \pm 15.8	39.0 \pm 5.2	0.064	R
<i>Expt. 3. July 1996</i>				
Carrizo	22.0 \pm 1.5	9.7 \pm 1.4	0.000	S
Swingle	29.8 \pm 2.3	9.1 \pm 1.4	0.000	S
HRS-1101	41.5 \pm 3.8	14.5 \pm 2.3	0.000	S
<i>Severinia buxifolia</i>	27.4 \pm 5.3	9.8 \pm 2.3	0.011	S
HRS-941	32.6 \pm 3.9	11.3 \pm 1.3	0.000	S
<i>Microcitrus australis</i>	33.0 \pm 5.9	9.7 \pm 1.5	0.002	S
<i>Murraya paniculata</i>	29.3 \pm 6.6	20.4 \pm 3.5	0.256	R
<i>Glycosmis pentaphylla</i>	37.5 \pm 5.2	32.9 \pm 4.1	0.501	R
<i>Eremocitrus glauca</i>	15.2 \pm 1.2	6.0 \pm 1.2	0.000	S
<i>Expt. 4. Dec. 1996</i>				
Carrizo	29.7 \pm 1.8	20.6 \pm 3.5	0.039	S
Swingle	41.0 \pm 3.7	20.0 \pm 4.5	0.003	S
HRS-802	66.3 \pm 5.5	44.3 \pm 5.5	0.015	S
HRS-812	48.7 \pm 5.1	27.1 \pm 2.0	0.002	S
HRS-896	41.7 \pm 4.3	29.5 \pm 3.2	0.043	S
Sun Chu Sha	62.8 \pm 9.2	45.8 \pm 4.4	0.120	R
<i>Triphasia trifolia</i>	26.4 \pm 3.4	11.5 \pm 1.0	0.001	S
<i>Clymenia polyandra</i>	29.2 \pm 7.0	5.4 \pm 1.3	0.006	S
<i>Citrus hystris</i>	17.8 \pm 5.0	5.6 \pm 1.7	0.037	S
<i>Fortunella hindsii</i>	27.2 \pm 8.6	4.0 \pm 0.5	0.055	R
<i>Expt. 5. Oct. 1997</i>				
Carrizo	26.4 \pm 2.8	9.2 \pm 1.2	0.000	S
Swingle	27.0 \pm 3.5	4.0 \pm 1.0	0.000	S
<i>Glycosmis pentaphylla</i>	23.1 \pm 3.2	7.7 \pm 1.0	0.001	S
<i>Expt. 6. May 1998</i>				
Carrizo	12.9 \pm 0.7	6.3 \pm 0.9	0.000	S
Swingle	18.3 \pm 1.8	7.3 \pm 1.0	0.000	S
<i>Glycosmis pentaphylla</i>	21.3 \pm 2.9	10.4 \pm 1.2	0.005	S
HRS-802	26.4 \pm 1.7	12.4 \pm 1.4	0.000	S
HRS-896	17.1 \pm 1.0	9.5 \pm 1.5	0.001	S
HRS-941	20.8 \pm 1.8	11.4 \pm 2.1	0.005	S
<i>Citrus aurantium</i>	20.7 \pm 1.1	4.2 \pm 0.5	0.000	S

¹For each species or cultivar, means for seven infested plants were compared with the mean of seven noninfested plants.

²Significance in *t* test.

³Ratings: S = susceptible; infested were significantly different from noninfested plants, $P < 0.05$. R = resistant; infested were not significantly different from noninfested plants, $P > 0.05$.

Table 4. Larval survival and growth on rootstocks in six experiments, means \pm se.

Species/cultivar	Larval survival		Larval growth	
	(%)	Rating	(mg)	Rating ¹
Expt. 1. Jan. 1996				
Carrizo	91 \pm 1	S	180 \pm 5	S
Swingle	94 \pm 2	S	151 \pm 6	R
Sun Chu Sha	94 \pm 3	S	173 \pm 4	S
HRS-801	76 \pm 2	R	151 \pm 8	R
HRS-802	89 \pm 5	S	206 \pm 4	S
HRS-812	93 \pm 4	S	160 \pm 7	S
HRS-896	87 \pm 4	S	138 \pm 8	HR
HRS-941	89 \pm 3	S	166 \pm 4	S
<i>Afraegle paniculata</i>	94 \pm 2	S	155 \pm 6	S
P (F test)	0.003		0.0001	
Expt. 2. June 1996				
Carrizo	96 \pm 3	S	185 \pm 8	S
Swingle	89 \pm 3	S	185 \pm 5	S
HRS-801	77 \pm 5	R	122 \pm 7	HR
HRS-802	96 \pm 2	S	196 \pm 11	S
HRS-896	87 \pm 4	S	158 \pm 5	S
Smooth Flat Seville	96 \pm 2	S	190 \pm 6	S
<i>Balsamocitrus dawei</i>	81 \pm 3	R	128 \pm 10	HR
Yuma	87 \pm 3	S	183 \pm 7	S
<i>Citropsis gilletiana</i>	96 \pm 3	S	258 \pm 12	HS
P (F test)	0.0001		0.0001	
Expt. 3. July 1996				
Carrizo	83 \pm 5	S	207 \pm 11	S
Swingle	86 \pm 5	S	213 \pm 20	S
HRS-1101	79 \pm 3	S	176 \pm 8	S
<i>Severinia buxifolia</i>	70 \pm 7	S	104 \pm 11	ffR
HRS-941	70 \pm 4	S	145 \pm 9	R
<i>Microcitrus australis</i>	63 \pm 7	S	80 \pm 6	HR
<i>Murraya paniculata</i>	87 \pm 4	S	164 \pm 12	S
<i>Glycosmis pentaphylla</i>	57 \pm 5	S	38 \pm 5	HR
<i>Eremocitrus glauca</i>	51 \pm 8	R	81 \pm 8	HR
P (F test)	0.0001		0.0001	
Expt. 4. Dec. 1996				
Carrizo	89 \pm 3	S	182 \pm 10	S
Swingle	87 \pm 4	S	171 \pm 10	S
HRS-802	93 \pm 3	S	228 \pm 5	HS
HRS-812	91 \pm 1	S	199 \pm 9	S
HRS-896	87 \pm 5	S	158 \pm 12	S
Sun Chu Sha	94 \pm 2	S	209 \pm 10	S
<i>Triphasia trifolia</i>	56 \pm 6	HR	55 \pm 4	HR
<i>Clymenia polyandra</i>	89 \pm 4	S	124 \pm 5	R
<i>Citrus hystrix</i>	83 \pm 2	S	89 \pm 12	HR
<i>Fortunella hindsii</i>	73 \pm 3	S	150 \pm 13	S
P (F test)	0.0001		0.0001	
Expt. 5. Oct. 1997				
Carrizo	86 \pm 4	S	203 \pm 14	S
Swingle	81 \pm 3	S	164 \pm 17	S
<i>Glycosmis pentaphylla</i>	67 \pm 7	R	30 \pm 5	HR
P (F test)	0.044		0.0001	
Expt. 6. May 1998				
Carrizo	81 \pm 7	S	156 \pm 9	S
Swingle	90 \pm 4	S	144 \pm 6	S
<i>Glycosmis pentaphylla</i>	61 \pm 7	S	35 \pm 3	HR
HRS-802	74 \pm 6	S	214 \pm 5	HS
HRS-896	79 \pm 6	S	116 \pm 9	R
HRS-941	80 \pm 4	S	145 \pm 9	S
<i>Citrus aurantium</i>	90 \pm 3	S	195 \pm 8	HS
P (F test)	0.012		0.0000	

¹Ratings: Carrizo is susceptible (S). All means rated as S were not distinguished from 'Carrizo' by Tukey's honestly significant difference (HSD) test at $P \leq 0.05$ and are also considered susceptible. HS = highly susceptible, more larval survival or growth than 'Carrizo', and distinguished from it by Tukey's HSD test at $P \leq 0.05$. R = resistant, less larval survival or growth than 'Carrizo', and distinguished from it by Tukey's HSD test at $P \leq 0.05$. HR = highly resistant, less larval survival or growth than 'Carrizo', and distinguished from it by Tukey's HSD test at $P \leq 0.001$.

Based on these experiments, the species *Balsamocitrus dawei* has a high level of resistance to *D. abbreviatus* larvae, suffering little root damage and inhibiting larval growth, while *G. pentaphylla*, *E. glauca*, *M. australis*, *S. buxifolia*, *T. trifolia*, and *C. hystrix* have a lesser level of resistance that was evidenced mainly by a depression in larval growth (Table 5). The cuttings of *C. hystrix* used in testing appeared healthy and similar to seedlings. However, it should be noted that if seedlings had been used for this cultivar, rather than cuttings, there would have been more genetic variability among the plants tested and differences in root structure might have affected test results. All of the commercially used rootstocks that we tested in these experiments ('Swingle', 'Smooth Flat Seville', 'Sun Chu Sha', and *C. aurantium*), many new hybrid rootstocks, and some of the related species were as susceptible to *D. abbreviatus* weevil larvae as was 'Carrizo'. One new hybrid rootstock, HRS-801, depressed the growth of *D. abbreviatus* larvae consistently in two experiments. Only *C. aurantium* was possibly more susceptible to *Diaprepes* than 'Carrizo', based on the resistance index.

Discussion

This series of experiments identified several sources of resistance to be used in future research and breeding efforts. Information in the literature on sexual and graft compatibility of citrus relatives is limited (Swingle, 1967). However, the species most resistant to *Diaprepes* weevil larvae, *Balsamocitrus dawei*, is considered sexually and graft incompatible with the genus *Citrus*, eliminating the opportunity to transmit the resistance factors by sexual hybridization or to use this species directly as a rootstock. The species *G. pentaphylla* is also sexually incompatible with citrus, but is partially graft-compatible with some citrus species (Bowman, unpublished data). A similar situation may exist with *T. trifolia*. Experiments are underway to identify suitable interstocks to allow use of *G. pentaphylla* rootstocks with commercial scions in areas infested by *D. abbreviatus*. The biochemical basis of resistance in *G. pentaphylla* has been investigated (Shapiro et al., 1997), and one component of this resistance has been identified as dehydrothalebanin (Shapiro et al., in press). This information might permit manipulation of biochemical pathways in existing citrus rootstocks, via genetic transformation, to induce "Glycosmis-type" resistance.

In contrast, antibiosis from *S. buxifolia*, *E. glauca*, *M. australis*, and *C. hystrix* could potentially be introgressed via breeding into citrus rootstocks because these four species are sexually compatible (Swingle, 1967; Medina-Filho et al., 1998) with some common species of the genus *Citrus*. Although resistance to *D. abbreviatus* was not a factor in their creation, hundreds of sexual hybrids between *Citrus* species and *Eremocitrus* or *Microcitrus* already exist in the germplasm of the Ft. Pierce USDA breeding programs.

BREEDING, CULTIVARS, ROOTSTOCKS, & GERMPLASM RESOURCES

Table 5. Resistance of citrus rootstocks and species to *D. abbreviatus* over six experiments.

Species/cultivar	Expts.	Root loss ^a (R1)	Larval growth ^a (R2)	Resistance rating ^a (R1+R2) ÷ 2
<i>Balsamocitrus dawei</i>	2	3	4	3.5
<i>Glycosmis pentaphylla</i>	3, 5, 6	2,3	4	3.2
<i>Eremocitrus glauca</i>	3	2	4	3
<i>Microcitrus australis</i>	3	2	4	3
<i>Severinia buxifolia</i>	3	2	4	3
<i>Triphasia trifolia</i>	4	2	4	3
<i>Citrus hystrix</i>	4	2	4	3
HRS-801	1, 2	2	3.5	2.8
<i>Murraya paniculata</i>	3	3	2	2.5
<i>Clymenia polyandra</i>	4	2	3	2.5
Smooth Flat Seville	2	3	2	2.5
Sun Chu Sha	1, 4	2.5	2	2.3
Hybrid HRS-896	1, 2, 4, 6	2	2.5	2.3
Hybrid HRS-941	1, 3, 6	2	2.3	2.2
Swingle	1, 2, 3, 4, 5, 6	2	2.2	2.1
Carrizo	1, 2, 3, 4, 5, 6	2	2	2
Yuma	2	2	2	2
HRS-812	1, 4	2	2	2
HRS-1101	3	2	2	2
<i>Afraegle paniculata</i>	1	2	2	2
<i>Citropsis gillettiana</i>	2	3	1	2
<i>Fortunella hindsii</i>	4	2	2	2
HRS-802	1, 2, 4, 6	2.3	1.5	1.9
<i>Citrus aurantium</i>	6	2	1	1.5

^aRatings: 1 = highly susceptible; 2 = susceptible; 3 = resistant; 4 = highly resistant.

Somatic hybrids between *Citrus* species and *Microcitrus* or *Severinia* have also been produced and are under evaluation (Grosser et al., 1996). Caution is still necessary in using these sources of resistance because we do not know how the short-term depression in larval growth that occurred from feeding on these four species relates to weevil maturation, reproduction, or population dynamics under field conditions. These species all suffered significant root damage from *D. abbreviatus* during the greenhouse testing.

The observation that hybrid HRS-801 significantly reduced larval growth and survival may indicate that a modest level of resistance occurs in some hybrids of *Citrus reticulata* x *Poncirus trifoliata*. This type of hybrid also has potential to combine other useful rootstock traits, including tree size control and resistance to citrus tristeza virus, citrus nematode, phytophthora diseases, cold, and citrus blight (Bowman, 1998; Wutscher and Bowman, 1999; Wutscher and Hill, 1995). However, we cannot eliminate the possibility that reduction in larval health for this selection may have been the result of a relatively small root mass (<25 g) for the tests with this selection, despite the conclusion that root weight (in the range of 25–45 g) was not associated with larval growth (Fig. 2). As with the antibiosis of *E. glauca*, *M. australis*, and *S. buxifolia*, and *C. hystrix*, better definition is needed for how this short-term depression in larval growth relates to longer-term resistance issues.

For future greenhouse testing, use of potting mix without perlite, as was done in

Expt. 6, may reduce the difficulty in recovering weevil larvae at the conclusion of tests. It did not appear to significantly affect testing of either standard susceptible cultivars like 'Carrizo' and 'Swingle' or the resistant species *G. pentaphylla*. Standardization of initial root mass of selections to be tested is important to minimize the potential influence of this factor on root damage and larval growth. *Glycosmis pentaphylla* appeared to be a good resistant control to include in future greenhouse tests for *Diaprepes* resistance; *B. dawei* may also be used for this purpose, but is probably less convenient because of a long juvenile period, greater seedling variability, and very limited seed sources in the United States.

The research described here has identified several new potential sources of resistance to *Diaprepes* weevil larvae, defined relative level of susceptibility for several common and new rootstock selections, and identified ways to improve greenhouse testing for weevil resistance. There appear to be several long-term approaches available for future development of new rootstocks resistant to *Diaprepes* weevil.

Literature Cited

Abrahamson, S.G. 1994. Attacking citrus root weevil, p. 10, 12, 15. In: Frank C. Garner (ed.), Florida Grower and Rancher.
 Beavers, J.B., R.E. Woodruff, S.A. Lovstrand, and W.J. Schroeder. 1979. Bibliography of the sugarcane rootstock borer weevil, *Diaprepes abbreviatus*. *Bul. Entomol. Soc. Amer.* 25:25–29.

Beavers, J.B. and D.J. Hutchison. 1985. Evaluation of selected *Citrus* spp. and relatives for susceptibility to root injury by *Diaprepes abbreviatus* larvae (Coleoptera: Curculionidae). *Fla. Entomol.* 68(1):222–223.
 Bowman, K.D. 1998. Performance of 'Fallglo' citrus hybrid on ten rootstocks in Lake County. *Proc. Fla. State Hort. Soc.* 111:177–180.
 Florida Dept. of Agriculture. 1998. Annual report. Bureau of citrus budwood registration. Div. Plant Industry. Winter Haven, Fla.
 Grosser, J.W. and C.W. McCoy. 1996. Feeding response of first instar larvae of *Diaprepes abbreviatus* to different novel intergeneric citrus somatic hybrids. *Proc. Fla. State Hort. Soc.* 109:62–66.
 Grosser, J.W., F.A.A. Mourao-Fo, F. G. Gmitter, Jr., E.S. Louzada, J. Jiang, K. Baergen, A. Quiros, C. Cabasson, J.L. Schell, and J.L. Chandler. 1996. Allotetraploid hybrids between citrus and seven related genera produced by somatic hybridization. *Theor. Appl. Genet.* 92:577–582.
 McCoy, C.W., L.W. Duncan, and E. D. Quintela. 1996. A review of IPM strategies for citrus root weevils with emphasis on microbial control. *Proc. Intl. Soc. Citricult.* 1996, Vol. 1:638–641.
 Medina-Filho, H.P., R. Bordignon, and R.M.L. Ballve. 1998. Sunkifolias and Buxisunkis: Sexually obtained reciprocal hybrids of *Citrus sunki* x *Severinia buxifolia*. *Genet. Mol. Biol.* 21:129–133.
 Norman, P.A., A.G. Selhime, and R.A. Sutton. 1974. Feeding damage to five citrus rootstocks by larvae of *Diaprepes abbreviatus* (Coleoptera: Curculionidae). *Fla. Entomol.* 57:296.
 Schroeder, W.J. 1988. Entomogenous nematodes for root weevil control in citrus, p. 1223–1226. In: R. Goren and K. Mendel (eds.), *Proc. Sixth Intl. Citrus Congr.*, Tel Aviv, Israel, 6–11 Mar., 1988. Balaban, Philadelphia.
 Schroeder, W.J. and J.B. Beavers. 1977. Citrus root weevils in Florida: identification, biology, and control. *Proc. Intl. Soc. Citricult.* 2:498–500.
 Shapiro, J.P., K.D. Bowman, and S.L. Lapointe. 2000. Dehydrothalebanin, a source of resistance from *Glycosmis pentaphylla* against the citrus root weevil *Diaprepes abbreviatus*. *J. Agr. Food Chem.* 48:4404–4409.
 Shapiro, J.P., K.D. Bowman, and H. Smith. 1997. Resistance of citrus rootstocks and *Glycosmis pentaphylla* against larval citrus root weevils, *Diaprepes abbreviatus*, in live root or diet-incorporation assays. *Fla. Entomol.* 80:471–477.
 Shapiro, J.P. and T.R. Gottwald. 1995. Resistance of eight cultivars of citrus rootstock to a larval root weevil, *Diaprepes abbreviatus* L. (Coleoptera: Curculionidae). *J. Econ. Entomol.* 88:148–154.
 Swingle, W.T. and P.C. Reece. 1967. The botany of citrus and its wild relatives, p. 190–430. In: W. Reuther, H.J. Webber, W.D. Batchelor (eds.), *The citrus industry*. Vol. 1. Univ. of Calif. Press, Berkeley.
 Wutscher, H.K. and K.D. Bowman. 1999. Performance of 'Valencia' orange on 21 rootstocks. *HortScience* 34:622–624.
 Wutscher, H.K. and L.L. Hill. 1995. Performance of 'Hamlin' orange on 16 rootstocks in east-central Florida. *HortScience* 30:41–43.

HORTSCIENCE

Inher Resis

James W. Department State Uni Gary G. Irrigated Universi

Additional genetics, n

Abstract. Pacific N caused by in the irri is know selection Research The obje resistanc quality, progenie leaf disk the prog orchard : 1 suscep derived

Pow mon in t ington b been es ered th sweet 1998: is a we ports. produ 'Van' to P. which devel may s and fruit

Rece for p of a men ingto Pehr and cha the Col the par reg ma 'T E-